



Faculty of Resource Science and Technology

**PHYTOCHEMICAL STUDIES AND PHARMACOPHORE ANALYSIS  
OF *SONNERATIA CASEOLARIS* FRUITS**

Liyana Zafirah Binti Othman

Bachelor of Science with Honours  
(Resource Chemistry)  
2016

# **Phytochemical Studies and Pharmacophore Analysis of *Sonneratia caseolaris*'s Fruits**

**Liyana Zafirah Binti Othman**

**(41916)**

This thesis is submitted in fulfillment of the requirement for the degree of  
Bachelor of Science with Honours  
(Resource Chemistry)

**Supervisor: Dr. Mohd Razip Bin Asaruddin**

Faculty of Resource Science and Technology  
University Malaysia Sarawak

2016

## ACKNOWLEDGMENT

Alhamdulillah, I am able to complete my thesis successfully. This thesis was able to be completed with all people that continuously support behind me.

First of all, I am very thankful to my family for continuous mentally, physically and financially support me all the time in completing this Final Year Project. I will not forget their countless prayer and endless efforts towards my entire journey in university. Next, I would like to express my sincere gratitude to my supervisor, Dr. Mohd. Razip Bin Asaruddin, for giving me the opportunity to conduct this research. Thank you for being a committed supervisor that helps me to understand well about my Final Year Project.

My appreciation also extends to our postgraduates students Mohammad Farhan Arifteen Rosli, Sam Ezekiel Radhakrishnan, Ardy Mursyid Romli, Racheal James and Tiara Nales Nyawai for devoted their time, monitoring and giving advices throughout my final year. I am grateful to have them who always share their knowledge and giving suggestions to overcome my obstacles during my research. And finally, I acknowledge all my lab mates and friends for their help, co-operation and willingness to accompany till midnight throughout the course of this work.

And most of all to God, who made me more determined to continue doing this work in spite of the obstacles and struggles that came. Thank you.

UNIVERSITI MALAYSIA SARAWAK

Grade: A

Please tick (✓)

Final Year Project Report

Masters

PhD

✓

DECLARATION OF ORIGINAL WORK

This declaration is made on the 14 day of JUNE 2016

Student's Declaration:

I Liyah Zafirah Binti Othman (41916) faculty of Resource Science and Technology  
(PLEASE INDICATE STUDENT'S NAME, MATRIC NO. AND FACULTY) hereby declare that the work entitled Phytochemical Studies and Pharmacophore Analysis of *Sonneratia caseolaris* Fruits is my original work. I have not copied from any other students' work or from any other sources except where due reference or acknowledgement is made explicitly in the text, nor has any part been written for me by another person.

14/6/2016

Date submitted

Liyah Zafirah Binti Othman (41916)

Name of the student (Matric No.)

Supervisor's Declaration:

I Dr. Mohd Razip Bin Asaruddin (SUPERVISOR'S NAME) hereby certifies that the work entitled Phytochemical Studies and Pharmacophore Analysis of *Sonneratia caseolaris* Fruit (TITLE) was prepared by the above named student, and was submitted to the "FACULTY" as a \* partial/full fulfillment for the conferment of Bachelor of Science with Honours (Resource Chemistry) (PLEASE INDICATE THE DEGREE), and the aforementioned work, to the best of my knowledge, is the said student's work.

Received for examination by: Dr. Mohd Razip Bin Asaruddin  
(Name of the supervisor)

Date: 14/6/2016



## TABLE OF CONTENTS

I declare that Project/Thesis is classified as (Please tick (✓)):

- ☐ **CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)\*  
☐ **RESTRICTED** (Contains restricted information as specified by the organisation where research was done)\*  
☒ **OPEN ACCESS**

### Validation of Project/Thesis

I therefore duly affirm with free consent and willingly declare that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abiding interest and rights as follows:

- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies for the purpose of academic and research only and not for other purpose.
- The Centre for Academic Information Services has the lawful right to digitalise the content for the Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic exchange between Higher Learning Institute.
- No dispute or any claim shall arise from the student itself neither third party on this Project/Thesis once it becomes the sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student except with UNIMAS permission.

Student signature \_\_\_\_\_

(Date) 14/6/2016

Supervisor signature: \_\_\_\_\_

(Date) 14/6/16

Current Address:

Lot 911 kampung Paya Jauh, 28000 Temerloh, Pahang Darul Makmur.

Notes: \* If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organisation with the period and reasons of confidentiality and restriction.

[The instrument is duly prepared by The Centre for Academic Information Services]

## TABLE OF CONTENTS

<b>ACKNOWLEDGEMENT.....</b>	<b>I</b>
<b>DECLARATION.....</b>	<b>II</b>
<b>TABLE OF CONTENTS.....</b>	<b>IV</b>
<b>LIST OF ABBREVIATION.....</b>	<b>VII</b>
<b>LIST OF FIGURES.....</b>	<b>IX</b>
<b>LIST OF TABLES.....</b>	<b>X</b>
<b>ABSTRACT/ ABSTRAK.....</b>	<b>1</b>
 <b>1.0 INTRODUCTION</b>	
1.1 General introduction .....	2
1.2 Objectives.....	4
 <b>2.0 LITERATURE REVIEW</b>	
2.1 Computer-aided drug design (CADD).....	5
2.1.1 Pharmacophore modelling .....	6
2.1.2 Structure-based drug design (SBDD) .....	7
2.1.3 Ligand-based drug design (LBDD) .....	8
2.2 <i>Sonneratia caseolaris</i> .....	8
2.3 Phytochemical constituents in <i>Sonneratia caseolaris</i> fruits .....	11
2.3.1 Flavanoids .....	11
2.3.2 Sterols .....	11
2.3.3 Triterpenoids .....	12
2.4 Medicinal values .....	14
2.4.1 Traditional uses .....	14
2.5 Biological activities .....	15
2.5.1 Antioxidant.....	15
2.5.2 Antimicrobial.....	15
2.5.3 Cytotoxicity .....	16
 <b>3.0 METHODOLOGY</b>	
3.1 Sample collection.....	17

3.2	General procedures .....	18
3.3	Sample preparation .....	20
3.4	Plant extraction .....	21
3.5	Isolation and purification of chemical constituents .....	22
3.5.1	Thin Layer Chromatography (TLC) .....	22
3.5.2	Column Chromatography (CC) .....	24
3.5.3	TLC detector.....	26
3.6	Structural elucidation of compound .....	27
3.6.1	Gas Chromatography-Mass Spectrometry (GC-MS) .....	27
3.6.2	Fourier Transform Infrared Spectrophotometer (FTIR) .....	27
3.6.3	Nuclear Magnetic Resonance (NMR) Spectroscopy .....	28
3.7	Pharmacophore modelling using CADD .....	30
3.7.1	Training set.....	30
3.7.2	Hypothesis generation.....	31
3.7.1	Virtual screening.....	32
3.8	Biological activity.....	34
3.8.1	Antimicrobial test .....	34
3.8.2	Cytotoxicity test.....	35
<b>4.0 RESULTS &amp; DISCUSSION</b>		
4.1	Extraction of <i>Sonneratia caseolaris</i> fruits .....	37
4.2	Isolation and purification of crude extract of <i>Sonneratia caseolaris</i> .....	38
4.2.1	Thin Layer Chromatography (TLC) analysis for methanol crude extracts.....	38
4.2.2	Column Chromatography of <i>S. caseolaris</i> .....	40
4.3	Structural Elucidation .....	45
4.3.1	Gas Chromatography-Mass Spectroscopy (GC-MS) .....	45
4.3.2	Nuclear Magnetic Resonance Spectroscopy (NMR) .....	47
4.3.3	Fourier Transform Infra-Red Spectrometer (FTIR) .....	49
4.3.4	Compound 1 .....	51
4.4	Pharmacophore modelling.....	52
4.4.1	Ligand-based drug design .....	52
4.5	Biological assay .....	60
4.5.1	Antimicrobial test .....	60
4.5.2	Cytotoxicity test.....	62

5.0 CONCLUSIONS .....	64
6.0 REFERENCES.....	66
7.0 APPENDIX	
A	A
B	B
C	C
D	D
E	E
F	F
G	G
H	H
I	I
J	J
K	K
L	L
M	M
N	N
O	O
P	P
Q	Q
R	R
S	S
T	T
U	U
V	V
W	W
X	X
Y	Y
Z	Z
AA	AA
AB	AB
AC	AC
AD	AD
AE	AE
AF	AF
AG	AG
AH	AH
AI	AI
AJ	AJ
AK	AK
AL	AL
AM	AM
AN	AN
AO	AO
AP	AP
AQ	AQ
AR	AR
AS	AS
AT	AT
AU	AU
AV	AV
AW	AW
AX	AX
AY	AY
AZ	AZ
BA	BA
BB	BB
BC	BC
BD	BD
BE	BE
BF	BF
BG	BG
BH	BH
BI	BI
BJ	BJ
BK	BK
BL	BL
BM	BM
BN	BN
BO	BO
BP	BP
BQ	BQ
BR	BR
BS	BS
BT	BT
BU	BU
BV	BV
BW	BW
BX	BX
BY	BY
BZ	BZ
CA	CA
CB	CB
CC	CC
CD	CD
CE	CE
CF	CF
CG	CG
CH	CH
CI	CI
CJ	CJ
CK	CK
CL	CL
CM	CM
CN	CN
CO	CO
CP	CP
CQ	CQ
CR	CR
CS	CS
CT	CT
CU	CU
CV	CV
CW	CW
CX	CX
CY	CY
CZ	CZ
DA	DA
DB	DB
DC	DC
DD	DD
DE	DE
DF	DF
DG	DG
DH	DH
DI	DI
DJ	DJ
DK	DK
DL	DL
DM	DM
DN	DN
DO	DO
DP	DP
DQ	DQ
DR	DR
DS	DS
DT	DT
DU	DU
DV	DV
DW	DW
DX	DX
DY	DY
DZ	DZ
EA	EA
EB	EB
EC	EC
ED	ED
EE	EE
EF	EF
EG	EG
EH	EH
EI	EI
EJ	EJ
EK	EK
EL	EL
EM	EM
EN	EN
EO	EO
EP	EP
EQ	EQ
ER	ER
ES	ES
ET	ET
EU	EU
EV	EV
EW	EW
EX	EX
EY	EY
EZ	EZ
FA	FA
FB	FB
FC	FC
FD	FD
FE	FE
FF	FF
FG	FG
FH	FH
FI	FI
FJ	FJ
FK	FK
FL	FL
FM	FM
FN	FN
FO	FO
FP	FP
FQ	FQ
FR	FR
FS	FS
FT	FT
FU	FU
FV	FV
FW	FW
FX	FX
FY	FY
FZ	FZ
GA	GA
GB	GB
GC	GC
GD	GD
GE	GE
GF	GF
GG	GG
GH	GH
GI	GI
GJ	GJ
GK	GK
GL	GL
GM	GM
GN	GN
GO	GO
GP	GP
GQ	GQ
GR	GR
GS	GS
GT	GT
GU	GU
GV	GV
GW	GW
GX	GX
GY	GY
GZ	GZ
HA	HA
HB	HB
HC	HC
HD	HD
HE	HE
HF	HF
HG	HG
HH	HH
HI	HI
HJ	HJ
HK	HK
HL	HL
HM	HM
HN	HN
HO	HO
HP	HP
HQ	HQ
HR	HR
HS	HS
HT	HT
HU	HU
HV	HV
HW	HW
HX	HX
HY	HY
HZ	HZ
IA	IA
IB	IB
IC	IC
ID	ID
IE	IE
IF	IF
IG	IG
IH	IH
II	II
IJ	IJ
IK	IK
IL	IL
IM	IM
IN	IN
IO	IO
IP	IP
IQ	IQ
IR	IR
IS	IS
IT	IT
IU	IU
IV	IV
IW	IW
IX	IX
IY	IY
IZ	IZ
JA	JA
JB	JB
JC	JC
JD	JD
JE	JE
JF	JF
JG	JG
JH	JH
JI	JI
JJ	JJ
JK	JK
JL	JL
JM	JM
JN	JN
JO	JO
JP	JP
JQ	JQ
JR	JR
JS	JS
JT	JT
JU	JU
JV	JV
JW	JW
JX	JX
JY	JY
JZ	JZ
KA	KA
KB	KB
KC	KC
KD	KD
KE	KE
KF	KF
KG	KG
KH	KH
KI	KI
KJ	KJ
KK	KK
KL	KL
KM	KM
KN	KN
KO	KO
KP	KP
KQ	KQ
KR	KR
KS	KS
KT	KT
KU	KU
KV	KV
KW	KW
KX	KX
KY	KY
KZ	KZ
LA	LA
LB	LB
LC	LC
LD	LD
LE	LE
LF	LF
LG	LG
LH	LH
LI	LI
LJ	LJ
LK	LK
LL	LL
LM	LM
LN	LN
LO	LO
LP	LP
LQ	LQ
LR	LR
LS	LS
LT	LT
LU	LU
LV	LV
LW	LW
LX	LX
LY	LY
LZ	LZ
MA	MA
MB	MB
MC	MC
MD	MD
ME	ME
MF	MF
MG	MG
MH	MH
MI	MI
MJ	MJ
MK	MK
ML	ML
MM	MM
MN	MN
MO	MO
MP	MP
MQ	MQ
MR	MR
MS	MS
MT	MT
MU	MU
MV	MV
MW	MW
MX	MX
MY	MY
MZ	MZ
NA	NA
NB	NB
NC	NC
ND	ND
NE	NE
NF	NF
NG	NG
NH	NH
NI	NI
NJ	NJ
NK	NK
NL	NL
NM	NM
NN	NN
NO	NO
NP	NP
NQ	NQ
NR	NR
NS	NS
NT	NT
NU	NU
NV	NV
NW	NW
NX	NX
NY	NY
NZ	NZ
OA	OA
OB	OB
OC	OC
OD	OD
OE	OE
OF	OF
OG	OG
OH	OH
OI	OI
OJ	OJ
OK	OK
OL	OL
OM	OM
ON	ON
OO	OO
OP	OP
OQ	OQ
OR	OR
OS	OS
OT	OT
OU	OU
OV	OV
OW	OW
OX	OX
OY	OY
OZ	OZ
PA	PA
PB	PB
PC	PC
PD	PD
PE	PE
PF	PF
PG	PG
PH	PH
PI	PI
PJ	PJ
PK	PK
PL	PL
PM	PM
PN	PN
PO	PO
PP	PP
PQ	PQ
PR	PR
PS	PS
PT	PT
PU	PU
PV	PV
PW	PW
PX	PX
PY	PY
PZ	PZ
QA	QA
QB	QB
QC	QC
QD	QD
QE	QE
QF	QF
QG	QG
QH	QH
QI	QI
QJ	QJ
QK	QK
QL	QL
QM	QM
QN	QN
QO	QO
QP	QP
QQ	QQ
QR	QR
QS	QS
QT	QT
QU	QU
QV	QV
QW	QW
QX	QX
QY	QY
QZ	QZ
RA	RA
RB	RB
RC	RC
RD	RD
RE	RE
RF	RF
RG	RG
RH	RH
RI	RI
RJ	RJ
RK	RK
RL	RL
RM	RM
RN	RN
RO	RO
RP	RP
RQ	RQ
RR	RR
RS	RS

## LIST OF ABBREVIATIONS

DCM	Dichloromethane
CHCl <sub>3</sub>	Chloroform
CD <sub>3</sub> OD	Deuterated methanol
EtOAc	Ethyl acetate
TLC	Thin Layer Chromatography
CC	Column Chromatography
R <sub>f</sub>	Retention factor
FT-IR	Fourier Transform Infrared Spectrophotometry
NMR	Nuclear Magnetic Resonance
<sup>1</sup> H-NMR	Proton Nuclear Magnetic Resonance
<sup>13</sup> C-NMR	Carbon Nuclear Magnetic Resonance
GC-MS	Gas Chromatography-Mass Spectrometry
GC-FID	Gas Chromatography-Flame Ionization Detector
UV-Vis	Ultraviolet Spectrophotometer
μL	microliter
LC <sub>50</sub>	Lethality concentration
Ppm	Part per million
mL	Milliliter
CADD	Computer-Aided Drug Design
SBDD	Structure-Based Drug Design
LBDD	Ligand-Based Drug Design
MeOH	Methanol
HBA	Hydrogen Bond Acceptor
HBD	Hydrogen Bond Donor
HI	Hydrophobic Interaction
AR	Aromatic Interaction
NIA	Negative Ionization Area

3-D	Three dimensional	88
<i>S. caseolaris</i>	<i>Sonneratia caseolaris</i>	
<i>E. coli</i>	<i>Escherichia coli</i>	
<i>S. aureus</i>	<i>Staphylococcus aureus</i>	
Figure 1.1	Diagram of chemical structures	11
Figure 1.2	Diagram of chemical structures	12
Figure 1.3	Diagram of chemical structures	13
Figure 1.4	The number of peaks collected (UNIMAS)	17
Figure 1.5	The first peak of the preparation	20
Figure 1.6	The first peak of the preparation	22
Figure 1.7	Diagram of the development of spots on TLC	23
Figure 1.8	Diagram of the development of spots on TLC	24
Figure 1.9	Diagram of the development of spots on TLC	25
Figure 1.10	Diagram of the development of spots on TLC	26
Figure 1.11	Diagram of the development of spots on TLC	28
Figure 1.12	Diagram of the development of spots on TLC	31
Figure 1.13	Diagram of the development of spots on TLC	32
Figure 1.14	Diagram of the development of spots on TLC	33
Figure 1.15	Diagram of the development of spots on TLC	34
Figure 1.16	Diagram of the development of spots on TLC	36
Figure 1.17	Diagram of the development of spots on TLC	39
Figure 1.18	Diagram of the development of spots on TLC	42
Figure 1.19	Diagram of the development of spots on TLC	43
Figure 1.20	Diagram of the development of spots on TLC	44
Figure 1.21	Diagram of the development of spots on TLC	45
Figure 1.22	Diagram of the development of spots on TLC	46
Figure 1.23	Diagram of the development of spots on TLC	47
Figure 1.24	Diagram of the development of spots on TLC	48
Figure 1.25	Diagram of the development of spots on TLC	50
Figure 1.26	Diagram of the development of spots on TLC	51
Figure 1.27	Diagram of the development of spots on TLC	52
Figure 1.28	Diagram of the development of spots on TLC	53
Figure 1.29	Diagram of the development of spots on TLC	54
Figure 1.30	Diagram of the development of spots on TLC	55
Figure 1.31	Diagram of the development of spots on TLC	56
Figure 1.32	Diagram of the development of spots on TLC	57
Figure 1.33	Diagram of the development of spots on TLC	58
Figure 1.34	Diagram of the development of spots on TLC	59
Figure 1.35	Diagram of the development of spots on TLC	60
Figure 1.36	Diagram of the development of spots on TLC	61
Figure 1.37	Diagram of the development of spots on TLC	62
Figure 1.38	Diagram of the development of spots on TLC	63
Figure 1.39	Diagram of the development of spots on TLC	64
Figure 1.40	Diagram of the development of spots on TLC	65
Figure 1.41	Diagram of the development of spots on TLC	66
Figure 1.42	Diagram of the development of spots on TLC	67
Figure 1.43	Diagram of the development of spots on TLC	68
Figure 1.44	Diagram of the development of spots on TLC	69
Figure 1.45	Diagram of the development of spots on TLC	70
Figure 1.46	Diagram of the development of spots on TLC	71
Figure 1.47	Diagram of the development of spots on TLC	72
Figure 1.48	Diagram of the development of spots on TLC	73
Figure 1.49	Diagram of the development of spots on TLC	74
Figure 1.50	Diagram of the development of spots on TLC	75
Figure 1.51	Diagram of the development of spots on TLC	76
Figure 1.52	Diagram of the development of spots on TLC	77
Figure 1.53	Diagram of the development of spots on TLC	78
Figure 1.54	Diagram of the development of spots on TLC	79
Figure 1.55	Diagram of the development of spots on TLC	80
Figure 1.56	Diagram of the development of spots on TLC	81
Figure 1.57	Diagram of the development of spots on TLC	82
Figure 1.58	Diagram of the development of spots on TLC	83
Figure 1.59	Diagram of the development of spots on TLC	84
Figure 1.60	Diagram of the development of spots on TLC	85
Figure 1.61	Diagram of the development of spots on TLC	86
Figure 1.62	Diagram of the development of spots on TLC	87
Figure 1.63	Diagram of the development of spots on TLC	88
Figure 1.64	Diagram of the development of spots on TLC	89
Figure 1.65	Diagram of the development of spots on TLC	90
Figure 1.66	Diagram of the development of spots on TLC	91
Figure 1.67	Diagram of the development of spots on TLC	92
Figure 1.68	Diagram of the development of spots on TLC	93
Figure 1.69	Diagram of the development of spots on TLC	94
Figure 1.70	Diagram of the development of spots on TLC	95
Figure 1.71	Diagram of the development of spots on TLC	96
Figure 1.72	Diagram of the development of spots on TLC	97
Figure 1.73	Diagram of the development of spots on TLC	98
Figure 1.74	Diagram of the development of spots on TLC	99
Figure 1.75	Diagram of the development of spots on TLC	100



## LIST OF FIGURES

Figures	Descriptions	Page
Figure 1.1	<i>Sonneratia caseolaris</i> plants	3
Figure 2.1	LigandScout 3.1 software	5
Figure 2.2	The workflow of pharmacophore modelling	6
Figure 2.3	The <i>S. caseolaris</i> fruit and leaves	10
Figure 2.4	Example of flavonoid structures	11
Figure 2.5	Example of sterol structures	12
Figure 2.6	Example of triterpenoid structures	13
Figure 3.1	The location for sample collected (UNIMAS)	17
Figure 3.2	The flow process of sample preparation	20
Figure 3.3	The flow process for plant extraction	22
Figure 3.4	Illustration for development of spots on TLC	23
Figure 3.5	General flow for CC	25
Figure 3.6	Column setup and the fraction collections	25
Figure 3.7	Vanillin stain solution	26
Figure 3.8	General flow for NMR Spectroscopy	28
Figure 3.9	The 3D structure of training sets	31
Figure 3.10	The training set and test set superimposed to each other	32
Figure 3.11	Example of pharmacophore model with the fit values	32
Figure 3.12	The flow process of pharmacophore modelling	33
Figure 3.13	The flow for antimicrobial test	34
Figure 3.14	The flow chart for cytotoxicity test	36
Figure 4.1	The best separation for methanol crude	39
Figure 4.2	TLC development of F <sub>5</sub> using CHCl <sub>3</sub> : EtOAc (1:2)	42
Figure 4.3	TLC spots for F <sub>5</sub> eluents	43
Figure 4.4	The dried fraction and TLC spot of compound 1	44
Figure 4.5	The result of GC-MS	45
Figure 4.6	The GC-MS results of each peak	46
Figure 4.7	The ion fragmentation for peak 10	46
Figure 4.8	The NMR result for compound 1	47
Figure 4.9	Zoom in of <sup>1</sup> H NMR spectrum result	48
Figure 4.10	FTIR spectrum for compound 1	50
Figure 4.11	Structure of compound 1	51
Figure 4.12	Pharmacophore Model generated from the training set	52
Figure 4.13	Interaction of Liposome with generated pharmacophore model	53
Figure 4.14	Interaction of Rifaximin with generated pharmacophore model	54
Figure 4.15	Interaction of Loperamide with generated pharmacophore model	54
Figure 4.16	Interaction of Linezolid with generated pharmacophore model	55
Figure 4.17	Interaction of luteolin with generated pharmacophore model	56
Figure 4.18	Interaction of lupeol with generated pharmacophore model	56
Figure 4.19	Interaction of compound 1 with generated pharmacophore model	57
Figure 4.20	Antimicrobial test for hexane crude	61
Figure 4.21	Antimicrobial test for methanol crude	61
Figure 4.22	Average death (%) of against log <sub>10</sub> concentration (µg/mL)	62

## LIST OF TABLES

Tables	Descriptions	Page
Table 2.1	Taxonomy of <i>Sonneratia caseolaris</i>	9
Table 4.1	The result for crude extractions	37
Table 4.2	Calculation of retention factor, R <sub>f</sub> value for compound observed in the methanol crude extract of <i>S. caseolaris</i> in the solvent system DCM: CH <sub>3</sub> Cl with ratio (1:9)	39
Table 4.3	Combined fractions from the first column	41
Table 4.4	The weight for combined fractions	42
Table 4.5	The combined fractions and the retention factor for F <sub>5</sub>	43
Table 4.6	Colour of the pharmacophore features	53
Table 4.7	Summary of pharmacophore feature analysis	58
Table 4.8	Zone of inhibition in diameter (mm)	60
Table 4.9	LC <sub>50</sub> determination for <i>Artemia salina</i>	62

# Phytochemical Studies and Pharmacophore Analysis of *Sonneratia caseolaris*'s Fruits

Liyana Zafirah Binti Othman

Department of Chemistry  
Faculty of Resource Science and Technology  
Universiti Malaysia Sarawak

## ABSTRACT

The aim for this study was to isolate the bioactive molecules, determine the biological activities and find the potential lead in *Sonneratia caseolaris* as antimicrobial drugs. Phytochemical examination on the methanol extract of the fruits gave about 37.37 g of crude and 4.82 % percentage yield which led to the isolation and characterization of compound 1. This structure compound were elucidated and characterized by spectroscopic analysis by using  $^1\text{H}$  and  $^{13}\text{C}$  Nuclear Magnetic Resonance (NMR) Spectroscopy, and Fourier Transform Infrared Spectroscopy (FTIR). In ligand-based pharmacophore modelling, the interaction of compound 1 with generated pharmacophore model shows the presence of hydrophobic (HY) and hydrogen bond acceptor (HBA) pharmacophore features with fit value of 35.4800. Biological evaluations were carried out on the methanol and hexane crude extracts. The results stated that the methanol crude extracts has higher  $\text{LC}_{50}$  value that was 2344.0 ppm. This higher value indicated that compound has no toxicity towards *Artemia salina*. In disc diffusion for antimicrobial test, it was found that *Sonneratia caseolaris* crude showed weak inhibition zone (7.33 mm and 7.00 mm) towards *Escherichia coli* and mild inhibition towards *Staphylococcus aureus* for both methanol and hexane crude.

Keywords: *Sonneratia caseolaris*, antimicrobial, pharmacophore modelling, toxicity, biological evaluations

## ABSTRAK

Tujuan kajian ini adalah untuk mengasingkan molekul bioaktif, menentukan aktiviti biologi dan mencari potensi dalam *Sonneratia caseolaris* sebagai ubat anti-mikrob. Kajian fitokimia atas ekstrak metanol daripada buah-buahan memberikan lebih kurang 37.37 g hasil peratusan mentah dan 4.82 % yang membawa kepada pengasingan dan pencirian sebatian 1. Sebatian struktur telah dijelaskan dan ciri-ciri analisis spektroskopi dengan menggunakan  $^1\text{H}$  dan  $^{13}\text{C}$  Nuclear Magnetic Resonance (NMR) Spektroskopi dan Fourier Transform Infrared Spektroskopi (FTIR). Berdasarkan ligan pemodelan pharmacophore, interaksi sebatian 1 dengan model pharmacophore dijana menunjukkan kehadiran hidrofobik (HY) dan ikatan hidrogen penerima (HBA) ciri-ciri pharmacophore dengan nilai patut 35.4800. Penilaian biologi telah dijalankan ke atas metanol dan ekstrak heksana mentah. Keputusan menyatakan bahawa ekstrak metanol mentah mempunyai nilai  $\text{LC}_{50}$  yang paling tinggi iaitu sebanyak 2344.0 ppm. Nilai ini lebih tinggi menunjukkan bahawa sebatian tidak toksik terhadap *Artemia salina*. Dalam penyebaran cakera untuk ujian antimikrob, didapati bahawa *Sonneratia caseolaris* mentah menunjukkan zon perencatan rendah (7.33 mm dan 7.00 mm) ke arah *Escherichia coli* dan perencatan sederhana ke arah *Staphylococcus aureus* untuk kedua-dua metanol dan heksana mentah.

Kata kunci: *Sonneratia caseolaris*, antimicrobial, pemodelan pharmacophore, toksik, penilaian biologi

## 1.0 INTRODUCTION

### 1.1 General introduction

Computer-aided Drug Design (CADD) is a computational methods and resources that are used to facilitate the design and discovery of new therapeutic solutions (Song *et al.*, 2009). It shows that search for new potent antimicrobial drug is essential to provide a safer treatment for the patients. A pharmacophore can be derived into two methods which are ligand-based and structure-based (Wobler *et al.*, 2008). However, it shows that there have been only a few studies on *Sonneratia caseolaris* regarding the pharmacological activities and discovery of new drug by using CADD. Therefore, this is an opportunity to develop and search for new potent antimicrobial drug from the fruit of *Sonneratia caseolaris* plant that could be commercialize.

*Sonneratia caseolaris* is one of typical mangrove tree species known as “kerala” that grows well in brackish water bodies in Sri Lanka (Abeywickrama & Jayasooriya, 2010) (Figure 1.1). In Malaysia, the fruit is known as “pedada” or “berembang”. *Sonneratia caseolaris* belongs to the family Sonneratiaceae (Nazli & Hashim, 2010). Howlader *et al.*, (2012) had states that *Sonneratia caseolaris* species is widespread and can be found easily in Asia. Based on previous research, *Sonneratia caseolaris* old fruit walls are used for helminthic infections, half-ripe fruit for coughs, while the pounded leaves for hematuria and small pox (Rahmatullah *et al.*, 2012). In addition, studies state that the calyces of *Sonneratia caseolaris* exhibited the strong antioxidant activity then followed by stamens (Bunyaphatsara *et al.*, 2003). As for toxicity test, it shows that the leaf of *Sonneratia caseolaris* methanolic was not toxic to *Artemia salina* (Melki *et al.*, 2011).



Previous studies states that steroids, flavonoids, triterpenoids and benzene carboxylic derivatives have been isolated and identified from the stems and twigs of *Sonneratia caseolaris* (Tian *et al.*, 2009). It shows that this plant contains high potential secondary metabolites that have medicinal value. The value of these secondary metabolites is increasing due to the constant discoveries of their potential role in drug development (Remani *et al.*, 2012). Figure 1.1 shows the plant that was discovered in this study.



**Figure 1.1** *Sonneratia caseolaris* plants

## 1.2 Objectives

This study will cover the following objectives:

1. To extract and isolate the chemical compounds from fruits of *S. caseolaris* by using chromatography techniques.
2. To characterize and identify the chemical constituents by using Gas Chromatography-Mass Spectrometry (GC-MS), Fourier Transform Infrared Spectroscopy (FTIR) and Nuclear Magnetic Resonance Spectroscopy (NMR).
3. To generate pharmacophore model by using commercial drugs.
4. To determine fit value of phytochemical constituents originate in *S. caseolaris* with commercial drugs by using the ligand-based pharmacophore modeling approach (LigandScout 3.1).
5. To determine the antimicrobial and cytotoxicity (*Artemia salina*) value of *S. caseolaris*'s fruit.



## 2.0 LITERATURE REVIEW

### 2.1 Computer-aided drug design (CADD)

Song *et al.*, (2009) states that Computer-aided drug design (CADD) is a specialized discipline that uses computational knowledge based methods to aid and facilitate the new drug discovery solutions. According to Kalyaanamoorthy & Chen, (2011) as for searching high affinity of flies repellent, a powerful molecular modelling technique has been useful in designing lead targets. To design good drugs, biological drug interactions should be considered such as using CADD to design a new drug of molecules in order to treat certain diseases in the human (McCarthy, 1999). The application of CADD was supported by using pharmacophore modelling. Chemical features based 3D pharmacophore models was built using computer software LigandScout 3.1 and used to evaluate and produced the geometrical structures of training set in their active forms (Figure 2.1). An established compound from Protein Data Bank (PDB) was used as training set. All structure of test set was generated and optimized using computer ChemDraw Ultra 12.0 software. Pharmacopore model were generated using a series of selected training set and then were mapped with test set. Figure 2.2 illustrated the work flow of pharmacophore modelling ~~that~~ will be used in this study.



**Figure 2.1** LigandScout 3.1 software




Figure 2.2 The workflow of pharmacophore modelling (Langer & Wolber, 2004)

### 2.1.1 Pharmacophore modelling

The pharmacophore can be derived both in a structure-based methods which between a ligand and its binding site, or in a ligand-based methods, which flexibly overlaying a set of active molecules and determining those conformations that are able to be overlapped in such a way that a maximum number of important chemical features geometrically overlap (Wobler *et al.*, 2008). Pharmacophore modeling approach is the technique rigidly models the interaction between a ligand and its binding site in a specific binding situation. The result is a three-dimensional (3D) spatial arrangement of chemical features. These feature maps, called 3D pharmacophores, can be used to search for similarities between binding situations or even for similarities between molecules.

Pharmacophore align molecule based on the three dimensional (3D) arrangement of chemical features. Noureen *et al.*, (2010), states that the generated pharmacophore hypothesis usually consist of five specific features which are one negatively ionizable group (NI), two hydrophobic aromatic (HY), one hydrogen bond acceptor (HBA) and one hydrogen bond donor (HBD).

Previous study on pharmacophore shows that the approach utilized in the study is successful in finding three potent Ftase inhibitors from Indian Plant Anticancer Database. Individually those three compounds are from group of vinca alkaloid & show lowest docked energy and hydrogen bonding stabilizes the interaction (Khan *et al.*, 2010). Besides, study shows that the discovery of ARRR.5 pharmacophore model was created by using indolizine derivatives acting as 15-LOX inhibitors. The hypothesis ARRR.5 might be considered as the best pharmacophore model because it displayed a good regression coefficient value of 0.9451, Pearson R 0.7439 and survival score 3.094 (Sharma & Kumar, 2015).

### **2.1.2 Structure-based drug design (SBDD)**

Structure-based drug design (SBDD) is the one with computational method shows that it assists to enhance progress in discovery and refinement of therapeutic agents (Marrone *et al.*, 1997). According to Kuntz, (1992) structure-based drug design refers to the complex process of using the information contained in the three-dimensional, 3D structure of a macromolecular target and related ligand-target complexes to design novel drugs for important human diseases. Structural information about a given macromolecular target will leads to a better

understanding of its specific function and enables the design of small molecule ligands that can bind to the target.

Pharmacophore modelling is one of the SBDD applications where it used training set (protein) from literature and database search (RCSB Protein Data Bank) to generate a pharmacophore model. This pharmacophore modelling was carried out using LigandScout 3.1 software.

### **2.1.3 Ligand-based drug design (LBDD)**

Ligand-based drug design is a method used in the absence of the receptor 3D information and it relies on knowledge of binding molecules to the biological target of interest. 3D quantitative structure activity relationships (3D QSAR) and pharmacophore modeling are the most important tools used in ligand-based drug design (Aparoy *et al.*, 2012).

## **2.2 *Sonneratia caseolaris***

*S. caseolaris* is one of typical mangrove tree species known as “kerala” that grows well in brackish water bodies in Sri Lanka (Abeywickrama & Jayasooriya, 2010). In Malaysia, the fruit is known as “pedada”, “berembang”, crabapple mangrove or mangrove apple. *S. caseolaris* belongs to the family Sonneratiaceae (Nazli & Hashim, 2010). *S. caseolaris* species is widespread and can be found in Bangladesh, Brunei Darussalam, Cambodia, China, India, Indonesia, Malaysia, Myanmar, Philippines, Singapore, Sri Lanka, Thailand, Viet Nam, Northeast Australia, Papua New Guinea, Solomon Islands, Vanuatu, New Caledonia, and Maldives (Md. S I. Howlader *et al.*, 2012). Table 2.1 shows the taxonomy of the *S. caseolaris* plants.



**Table 2.1** Taxonomy of *Sonneratia caseolaris*

<b>Kingdom</b>	<b>Plantae (Plants)</b>
<b>Division</b>	Tracheophyta
<b>Class</b>	Magnoliopsida
<b>Series</b>	Myrtales
<b>Family</b>	Sonneratiaceae
<b>Genus</b>	<i>Sonneratia</i>
<b>Species</b>	<i>Sonneratia caseolaris</i>



**Figure 2.3** The *S. caseolaris* fruit and leaves

Ahmed *et al.*, (2010) has states that the mangrove tree has oblong or obovate-elliptic coriaceous leaves and large red flowers. The tree also can grow up to 15-20 meters high. This typical non-viviporous mangrove species have their own benefits such as the fruit is edible, sap is used as a skin cosmetic, and leaves are used for feeding goats (Avenido & Serrano, 2012) (Figure 2.3). The ripened fruits of *S. caseolaris* have an appealing flavor and taste that can be used to prepare a delicious fruit drink. The fruit also contains a large number of small

seeds embedded in the fleshy part which are bitter if eaten (Abeywickrama & Jayasooriya, 2010). The tree flowers are edible as vegetable with *nampriks* (spicy dish). “Pedada” is a sour tasting young berry fruits that are edible (Wetwitayaklung *et al.*, 2013). In addition, the mature fruits really have a cheese like taste and can be eaten raw or cooked. Its stem and branch are used for firewood, building boats, posts of bridges and houses. Other than that, it can absorb, accumulate, distribute and circulate that heavy metal such as Cu, Pb, Zn, Cr and Ni in mangrove community (Wetwitayaklung *et al.*, 2013).



## 2.3 Phytochemical constituents in *Sonneratia caseolaris* fruits

### 2.3.1 Flavonoids

Flavonoids is one of secondary metabolites group which responsible for antioxidant activity. Based on previous studies, two flavonoids such as luteolin (Figure 2.4a) and luteolin-7-O- $\beta$ -glucoside (cynaroside) (Figure 2.4b) have been isolated from *Sonneratia caseolaris* leaves (Wetwitayaklung *et al.*, 2013).





Figure 2.4 Example of flavonoid structures

### 2.3.2 Sterols

Sterols are also known as steroid alcohols which occur naturally in plants. Based on previous study, sterols compound have been identified from the stem and twigs of *S. caseolaris*. The sterols that have been isolated are 6'-O-Acetyl- $\beta$ -daucosterol,  $\beta$ -Sitosterol (Figure 2.5a), Daucosterol (Figure 2.5c),  $\beta$ -Sitosterol palmitate (Figure 2.5d), Stigmast-5-en-3 $\beta$ -O-(6-O-hexadecanoyl- $\beta$ -D-glucopyranoside, and Cholest-5-en-3 $\beta$ ,7 $\alpha$ -diol (Tian *et al.*, 2009). After that, Tiwari *et al.*, (2010) states that through extraction and isolation, bioactive compound from the methanol extracts of pedada fruit such as  $\beta$ -sitosterol- $\beta$ -D-glucopyranoside were identified.



**Figure 2.5** Example of sterol structures

### 2.3.3 Triterpenoids

Triterpenoids are one of the secondary metabolites that represent the largest group of phytochemicals and research shown that it can be found in various plants including fruits and medicinal herbs (Bishayee *et al.*, 2011). Previous study shows that maslinic acid, betulin, lupeol and oleanolic can be found in *S. caseolaris* plants (Figure 2.6). Besides, oleanolic acid is well known to possess anti-inflammatory, antihyperlipidemic and hepatoprotective activities and sold as oral drug for human liver disorder in China (Liu, 1995) (Figure 2.6d).